

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(*E,E*)-1-Methyl-2,6-distyrylpyridinium iodideNarsimha Reddy Penthala,^a Joshua Eldridge,^a
Thirupathi Reddy Yerram Reddy,^a Sean Parkin^b and
Peter A. Crooks^{a*}^aDepartment of Pharmaceutical Sciences, College of Pharmacy, University of Kentucky, Lexington, KY 40536, USA, and ^bDepartment of Chemistry, University of Kentucky, Lexington, KY 40506, USA
Correspondence e-mail: pcrooks@uky.edu


Received 3 March 2010; accepted 18 May 2010

Key indicators: single-crystal X-ray study; $T = 90$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.021; wR factor = 0.055; data-to-parameter ratio = 15.1.

In the title compound, $\text{C}_{22}\text{H}_{20}\text{N}^+\cdot\text{I}^-$, the dihedral angles between the central pyridine ring and two outer benzene rings are 15.30 (10) and 11.82 (11)°. There are intermolecular π - π stacking interactions between the nearest phenyl ring over an inversion-related pyridyl ring, the shortest centroid-centroid distance being 3.672 (3) Å. The crystal structure of the compound indicates the 2,6-distyryl substituents have an *E* configuration.

Related literature

For the conventional synthesis, see: Stanek *et al.* (1952). For the activity of related compounds, see Zheng *et al.* (2005).



Experimental

Crystal data

$\text{C}_{22}\text{H}_{20}\text{N}^+\cdot\text{I}^-$
 $M_r = 425.29$
 Monoclinic, $C2/c$
 $a = 18.7309$ (4) Å
 $b = 9.5687$ (2) Å
 $c = 19.9829$ (4) Å
 $\beta = 90.279$ (1)°

$V = 3581.50$ (13) Å³
 $Z = 8$
 Cu $K\alpha$ radiation
 $\mu = 14.04$ mm⁻¹
 $T = 90$ K
 $0.18 \times 0.14 \times 0.08$ mm

Data collection

Bruker X8 Proteum diffractometer
 Absorption correction: multi-scan
 (SADABS; Bruker, 2001)
 $T_{\min} = 0.211$, $T_{\max} = 0.400$

24972 measured reflections
 3303 independent reflections
 3286 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.043$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.021$
 $wR(F^2) = 0.055$
 $S = 1.09$
 3303 reflections

219 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.58$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.49$ e Å⁻³

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and local procedures.

This investigation was supported by the NSF MRI grant CHE 0319176 (to SP). We thank Dr Greg Elliott for providing the microwave facilities.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NK2028).

References

- Bruker (2001). SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2007). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Stanek, J., Hebky, J. & Zverina, V. (1952). *Chem. Listy*, **46**, 735–736.
 Zheng, G., Dwoskin, L. P., Deaciuc, A. G., Norrholm, S. D. & Crooks, P. A. (2005). *J. Med. Chem.* **48**, 5551–5560.

supplementary materials

Acta Cryst. (2010). E66, o1793 [doi:10.1107/S1600536810018519]

(*E,E*)-1-Methyl-2,6-distyrylpyridinium iodide

N. R. Penthala, J. Eldridge, T. R. Y. Reddy, S. Parkin and P. A. Crooks

Comment

In continuation of our work on the lobeline analogues, and structure-affinity relationships of novel ligands for the vesicular monoamine transporter (Zheng *et al.*, 2005), we have undertaken the design, synthesis and structural analysis of a series of phenyl substituted *N*-alkyl distyrylpyridine analogs. The primary reasons for the X-ray analysis of the title compound was to confirm the geometry of 2,6-distyryl groups and to obtain detailed information on the molecular structure. This information will be useful in structure-activity relationship (SAR) studies. The title compound was prepared by the reaction of *N*-methyl, 2,6-lutidine iodide with benzaldehyde in the presence of pyrrolidine in ethanol under microwave irradiation at 25-29 W power level and at 130 °C for 3 minutes (Biotage microwave initiator). The compound was recrystallized from ethanol. The molecular structure and the atom-numbering scheme are shown in Fig.1. The X-ray studies revealed that the 2,6-distyryl substituents in the title compound both have *E* geometry. The central pyridine ring makes a dihedral angle of 15.30 (10)° and 11.82 (11)° with the adjacent phenyl rings.

Experimental

A mixture of *N*-methyl, 2,6-lutidine iodide (0.249 g, 1.0 mmol), benzaldehyde (0.300 g, 2.5 mmol) and pyrrolidine (6 µl) in ethanol (2 ml) was placed in a microwave-ready pressure vial equipped with a stirbar and irradiated in a Biotage microwave initiator for 3 minutes with the temperature set at 130 °C, at a power range of 25-29 W at 5 bar pressure. The cooled reaction mixture was taken out of the initiator, diluted with ethyl acetate, and filtered to afford a crude yellow solid. Crystallization from alcohol produced a yellow crystalline product of *N*-methyl-2,6-(*E*)distyrylpyridinium iodide that was suitable for X-ray analysis. ¹H NMR (DMSO *d*₆): δ 4.28 (*s*, 3H), 7.47-7.49 (*d*, *J*=6 Hz, 4H), 7.62-7.27 (*dd*, *J*=30.3 Hz, *J*=15.9 Hz, 6H), 7.848-7.872 (*d*, *J*=7.2 Hz, 4H), 8.26-8.29 (*m*, 2H), 8.39-8.44 (*t*, *J*=7.8 Hz, 1H); ¹³C NMR (DMSO *d*₆): δ 41.97, 119.09, 124.02, 128.40, 128.91, 130.39, 134.87, 142.22, 143.11, 149.88, 153.10.

Refinement

H atoms were found in difference Fourier maps and subsequently placed in idealized positions with constrained distances of 0.98 Å (RCH₃), 0.95 Å (C_{Ar}H), and with *U*_{iso}(H) values set to either 1.2*U*_{eq} or 1.5*U*_{eq} (RCH₃) of the attached atom.

Figures




Fig. 1. A view of the molecule with the atom numbering scheme. Displacement ellipsoids are drawn at the 50% probability level.

(*E,E*)-1-Methyl-2,6-distyrylpyridinium iodide

Crystal data

$C_{22}H_{20}N^+I^-$	$F(000) = 1696$
$M_r = 425.29$	$D_x = 1.577 \text{ Mg m}^{-3}$
Monoclinic, $C2/c$	Cu $K\alpha$ radiation, $\lambda = 1.54178 \text{ \AA}$
Hall symbol: $-C 2yc$	Cell parameters from 9883 reflections
$a = 18.7309 (4) \text{ \AA}$	$\theta = 4.4\text{--}68.5^\circ$
$b = 9.5687 (2) \text{ \AA}$	$\mu = 14.04 \text{ mm}^{-1}$
$c = 19.9829 (4) \text{ \AA}$	$T = 90 \text{ K}$
$\beta = 90.279 (1)^\circ$	Shard, yellow
$V = 3581.50 (13) \text{ \AA}^3$	$0.18 \times 0.14 \times 0.08 \text{ mm}$
$Z = 8$	

Data collection

Bruker X8 Proteum diffractometer	3303 independent reflections
Radiation source: fine-focus rotating anode graded multilayer optics	3286 reflections with $I > 2\sigma(I)$
Detector resolution: $5.6 \text{ pixels mm}^{-1}$	$R_{\text{int}} = 0.043$
φ and ω scans	$\theta_{\text{max}} = 68.5^\circ$, $\theta_{\text{min}} = 4.4^\circ$
Absorption correction: multi-scan (<i>SADABS</i> in <i>APEX2</i> ; Bruker, 2001)	$h = -22 \rightarrow 22$
$T_{\text{min}} = 0.211$, $T_{\text{max}} = 0.400$	$k = -11 \rightarrow 11$
24972 measured reflections	$l = -24 \rightarrow 24$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.021$	H-atom parameters constrained
$wR(F^2) = 0.055$	$w = 1/[\sigma^2(F_o^2) + (0.0309P)^2 + 5.1348P]$
$S = 1.09$	where $P = (F_o^2 + 2F_c^2)/3$
3303 reflections	$(\Delta/\sigma)_{\text{max}} = 0.003$
219 parameters	$\Delta\rho_{\text{max}} = 0.58 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.49 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: <i>SHELXTL</i> (Sheldrick, 2008), $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$
	Extinction coefficient: 0.000067 (15)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
I1	0.157566 (6)	0.613743 (12)	0.396017 (5)	0.01533 (8)
N1	0.38671 (8)	0.46130 (16)	0.46816 (7)	0.0115 (3)
C1	0.36127 (12)	0.60507 (19)	0.45527 (12)	0.0192 (5)
H1A	0.4014	0.6634	0.4410	0.029*
H1B	0.3407	0.6437	0.4963	0.029*
H1C	0.3248	0.6034	0.4199	0.029*
C2	0.44830 (10)	0.4136 (2)	0.43884 (9)	0.0133 (4)
C3	0.47346 (10)	0.2818 (2)	0.45649 (9)	0.0153 (4)
H3	0.5164	0.2477	0.4374	0.018*
C4	0.43651 (10)	0.2005 (2)	0.50154 (9)	0.0146 (4)
H4	0.4544	0.1112	0.5139	0.018*
C5	0.37368 (10)	0.24873 (19)	0.52856 (9)	0.0136 (4)
H5	0.3477	0.1915	0.5588	0.016*
C6	0.34807 (11)	0.37993 (18)	0.51204 (10)	0.0120 (4)
C7	0.28002 (10)	0.4316 (2)	0.53757 (9)	0.0133 (4)
H7	0.2554	0.5026	0.5135	0.016*
C8	0.25134 (11)	0.38157 (18)	0.59407 (10)	0.0146 (4)
H8	0.2794	0.3177	0.6194	0.018*
C9	0.18083 (10)	0.4165 (2)	0.62012 (9)	0.0139 (4)
C10	0.15174 (11)	0.3314 (2)	0.67007 (10)	0.0177 (4)
H10	0.1795	0.2576	0.6884	0.021*
C11	0.08281 (11)	0.3536 (2)	0.69312 (10)	0.0191 (4)
H11	0.0636	0.2945	0.7267	0.023*
C12	0.04199 (10)	0.4616 (2)	0.66730 (10)	0.0180 (4)
H12	-0.0055	0.4758	0.6823	0.022*
C13	0.07150 (11)	0.5492 (2)	0.61904 (9)	0.0169 (4)
H13	0.0442	0.6250	0.6021	0.020*
C14	0.13969 (10)	0.5276 (2)	0.59548 (9)	0.0154 (4)
H14	0.1588	0.5882	0.5625	0.018*
C15	0.48534 (11)	0.5004 (2)	0.38947 (10)	0.0189 (4)
H15	0.4637	0.5858	0.3760	0.023*
C16	0.54716 (11)	0.4660 (2)	0.36280 (10)	0.0183 (4)
H16	0.5664	0.3784	0.3761	0.022*

supplementary materials

C17	0.58969 (10)	0.5469 (2)	0.31504 (9)	0.0150 (4)
C18	0.64612 (11)	0.4793 (2)	0.28280 (10)	0.0168 (4)
H18	0.6546	0.3828	0.2909	0.020*
C19	0.68995 (11)	0.5521 (2)	0.23897 (10)	0.0188 (4)
H19	0.7273	0.5047	0.2163	0.023*
C20	0.67921 (11)	0.6936 (2)	0.22840 (10)	0.0190 (4)
H20	0.7094	0.7435	0.1987	0.023*
C21	0.62407 (11)	0.7629 (2)	0.26133 (10)	0.0209 (4)
H21	0.6173	0.8603	0.2547	0.025*
C22	0.57920 (11)	0.6895 (2)	0.30369 (10)	0.0171 (4)
H22	0.5410	0.7366	0.3252	0.020*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
I1	0.01676 (10)	0.01507 (10)	0.01419 (10)	0.00358 (4)	0.00364 (6)	0.00323 (4)
N1	0.0131 (7)	0.0105 (7)	0.0109 (7)	0.0004 (6)	0.0008 (6)	-0.0004 (6)
C1	0.0188 (11)	0.0122 (10)	0.0267 (12)	0.0042 (7)	0.0086 (9)	0.0059 (7)
C2	0.0133 (9)	0.0154 (9)	0.0113 (9)	0.0005 (7)	0.0013 (7)	-0.0029 (7)
C3	0.0151 (9)	0.0161 (9)	0.0148 (9)	0.0033 (7)	0.0025 (7)	-0.0012 (7)
C4	0.0179 (9)	0.0114 (8)	0.0145 (9)	0.0010 (7)	-0.0013 (7)	-0.0004 (7)
C5	0.0169 (9)	0.0126 (9)	0.0112 (8)	-0.0013 (7)	0.0007 (7)	-0.0001 (7)
C6	0.0133 (10)	0.0146 (9)	0.0080 (9)	-0.0014 (6)	0.0005 (7)	-0.0011 (6)
C7	0.0144 (9)	0.0119 (9)	0.0134 (9)	0.0014 (7)	0.0010 (7)	-0.0005 (7)
C8	0.0148 (10)	0.0155 (10)	0.0136 (10)	0.0006 (7)	0.0004 (8)	0.0013 (6)
C9	0.0154 (9)	0.0166 (9)	0.0098 (9)	-0.0011 (8)	0.0006 (7)	-0.0006 (7)
C10	0.0181 (10)	0.0192 (10)	0.0156 (9)	0.0010 (8)	0.0013 (7)	0.0041 (8)
C11	0.0191 (10)	0.0238 (10)	0.0145 (9)	-0.0037 (9)	0.0046 (8)	0.0016 (8)
C12	0.0150 (9)	0.0234 (10)	0.0156 (9)	-0.0009 (8)	0.0028 (7)	-0.0048 (8)
C13	0.0195 (9)	0.0175 (9)	0.0137 (9)	0.0047 (8)	0.0009 (7)	-0.0020 (7)
C14	0.0200 (10)	0.0156 (9)	0.0106 (9)	-0.0003 (8)	0.0029 (7)	-0.0003 (7)
C15	0.0223 (10)	0.0150 (9)	0.0193 (10)	0.0041 (8)	0.0063 (8)	0.0037 (8)
C16	0.0204 (10)	0.0139 (9)	0.0208 (10)	0.0013 (8)	0.0062 (8)	0.0020 (8)
C17	0.0152 (9)	0.0181 (10)	0.0117 (9)	-0.0004 (8)	0.0008 (7)	-0.0001 (7)
C18	0.0194 (10)	0.0149 (9)	0.0161 (9)	0.0012 (8)	0.0036 (7)	-0.0002 (7)
C19	0.0190 (10)	0.0222 (10)	0.0154 (9)	-0.0005 (8)	0.0056 (8)	-0.0016 (8)
C20	0.0212 (10)	0.0224 (10)	0.0133 (9)	-0.0019 (8)	0.0042 (7)	0.0035 (8)
C21	0.0257 (11)	0.0187 (10)	0.0184 (10)	0.0025 (8)	0.0022 (8)	0.0049 (8)
C22	0.0177 (9)	0.0184 (10)	0.0151 (9)	0.0042 (8)	0.0031 (7)	0.0018 (7)

Geometric parameters (\AA , $^\circ$)

N1—C2	1.374 (2)	C11—C12	1.384 (3)
N1—C6	1.380 (2)	C11—H11	0.9500
N1—C1	1.478 (2)	C12—C13	1.394 (3)
C1—H1A	0.9800	C12—H12	0.9500
C1—H1B	0.9800	C13—C14	1.379 (3)
C1—H1C	0.9800	C13—H13	0.9500
C2—C3	1.391 (3)	C14—H14	0.9500

C2—C15	1.467 (3)	C15—C16	1.319 (3)
C3—C4	1.379 (3)	C15—H15	0.9500
C3—H3	0.9500	C16—C17	1.467 (3)
C4—C5	1.377 (3)	C16—H16	0.9500
C4—H4	0.9500	C17—C22	1.397 (3)
C5—C6	1.383 (3)	C17—C18	1.399 (3)
C5—H5	0.9500	C18—C19	1.391 (3)
C6—C7	1.461 (3)	C18—H18	0.9500
C7—C8	1.341 (3)	C19—C20	1.385 (3)
C7—H7	0.9500	C19—H19	0.9500
C8—C9	1.461 (3)	C20—C21	1.395 (3)
C8—H8	0.9500	C20—H20	0.9500
C9—C10	1.401 (3)	C21—C22	1.387 (3)
C9—C14	1.401 (3)	C21—H21	0.9500
C10—C11	1.389 (3)	C22—H22	0.9500
C10—H10	0.9500		
C2—N1—C6	121.83 (16)	C12—C11—C10	120.22 (19)
C2—N1—C1	120.35 (16)	C12—C11—H11	119.9
C6—N1—C1	117.78 (16)	C10—C11—H11	119.9
N1—C1—H1A	109.5	C11—C12—C13	119.12 (18)
N1—C1—H1B	109.5	C11—C12—H12	120.4
H1A—C1—H1B	109.5	C13—C12—H12	120.4
N1—C1—H1C	109.5	C14—C13—C12	121.16 (18)
H1A—C1—H1C	109.5	C14—C13—H13	119.4
H1B—C1—H1C	109.5	C12—C13—H13	119.4
N1—C2—C3	118.49 (17)	C13—C14—C9	120.16 (18)
N1—C2—C15	119.95 (17)	C13—C14—H14	119.9
C3—C2—C15	121.55 (17)	C9—C14—H14	119.9
C4—C3—C2	120.45 (18)	C16—C15—C2	123.32 (18)
C4—C3—H3	119.8	C16—C15—H15	118.3
C2—C3—H3	119.8	C2—C15—H15	118.3
C5—C4—C3	119.95 (18)	C15—C16—C17	127.73 (19)
C5—C4—H4	120.0	C15—C16—H16	116.1
C3—C4—H4	120.0	C17—C16—H16	116.1
C4—C5—C6	120.46 (17)	C22—C17—C18	118.89 (17)
C4—C5—H5	119.8	C22—C17—C16	122.98 (17)
C6—C5—H5	119.8	C18—C17—C16	118.00 (18)
N1—C6—C5	118.76 (17)	C19—C18—C17	120.53 (19)
N1—C6—C7	119.48 (16)	C19—C18—H18	119.7
C5—C6—C7	121.71 (17)	C17—C18—H18	119.7
C8—C7—C6	121.80 (18)	C20—C19—C18	120.00 (18)
C8—C7—H7	119.1	C20—C19—H19	120.0
C6—C7—H7	119.1	C18—C19—H19	120.0
C7—C8—C9	125.79 (18)	C19—C20—C21	120.00 (18)
C7—C8—H8	117.1	C19—C20—H20	120.0
C9—C8—H8	117.1	C21—C20—H20	120.0
C10—C9—C14	118.44 (18)	C22—C21—C20	119.98 (19)
C10—C9—C8	118.45 (18)	C22—C21—H21	120.0
C14—C9—C8	123.05 (17)	C20—C21—H21	120.0

supplementary materials

C11—C10—C9	120.86 (19)	C21—C22—C17	120.56 (18)
C11—C10—H10	119.6	C21—C22—H22	119.7
C9—C10—H10	119.6	C17—C22—H22	119.7
C6—N1—C2—C3	-2.8 (3)	C8—C9—C10—C11	-175.20 (18)
C1—N1—C2—C3	174.87 (18)	C9—C10—C11—C12	-0.6 (3)
C6—N1—C2—C15	176.36 (17)	C10—C11—C12—C13	-1.3 (3)
C1—N1—C2—C15	-6.0 (3)	C11—C12—C13—C14	1.8 (3)
N1—C2—C3—C4	1.1 (3)	C12—C13—C14—C9	-0.2 (3)
C15—C2—C3—C4	-178.04 (18)	C10—C9—C14—C13	-1.7 (3)
C2—C3—C4—C5	1.0 (3)	C8—C9—C14—C13	175.49 (18)
C3—C4—C5—C6	-1.5 (3)	N1—C2—C15—C16	175.08 (19)
C2—N1—C6—C5	2.3 (3)	C3—C2—C15—C16	-5.8 (3)
C1—N1—C6—C5	-175.42 (18)	C2—C15—C16—C17	-177.77 (19)
C2—N1—C6—C7	-175.13 (17)	C15—C16—C17—C22	17.0 (3)
C1—N1—C6—C7	7.1 (3)	C15—C16—C17—C18	-167.2 (2)
C4—C5—C6—N1	-0.1 (3)	C22—C17—C18—C19	-1.5 (3)
C4—C5—C6—C7	177.27 (18)	C16—C17—C18—C19	-177.53 (18)
N1—C6—C7—C8	-158.92 (18)	C17—C18—C19—C20	1.9 (3)
C5—C6—C7—C8	23.7 (3)	C18—C19—C20—C21	-0.5 (3)
C6—C7—C8—C9	-173.77 (18)	C19—C20—C21—C22	-1.2 (3)
C7—C8—C9—C10	164.66 (19)	C20—C21—C22—C17	1.5 (3)
C7—C8—C9—C14	-12.5 (3)	C18—C17—C22—C21	-0.2 (3)
C14—C9—C10—C11	2.1 (3)	C16—C17—C22—C21	175.62 (19)

Fig. 1

