

Bis[tris(1*H*-pyrazol-1-yl- κ N²)methane]-nickel(II) bis[[tris(1*H*-pyrazol-1-yl- κ N²)methane]tris(thiocyanato- κ N)-nickelate(II)] methanol disolvate

Ganna Lyubartseva,^{a*} Sean Parkin^b and Uma Prasad Mallik^a

^aDepartment of Chemistry and Physics, Southern Arkansas University, Magnolia, AR 71753, USA, and ^bDepartment of Chemistry, University of Kentucky, Lexington, KY 40506, USA

Correspondence e-mail: gannalyubartseva@saumag.edu


Received 20 October 2011; accepted 27 October 2011

Key indicators: single-crystal X-ray study; $T = 90$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; disorder in main residue; R factor = 0.049; wR factor = 0.128; data-to-parameter ratio = 13.4.

Attempts to prepare the mononuclear $[(\text{tpm})\text{Ni}^{\text{II}}L_3]^{-1}$ [$\text{tpm} = \text{tris}(1H\text{-pyrazol-1-yl})\text{methane}$ and $L = \text{thiocyanate}$] anion yielded the methanol-solvated salt, $[(\text{tpm})_2\text{Ni}^{\text{II}}][(\text{tpm})\text{Ni}^{\text{II}}(\text{NCS})_3] \cdot 2\text{CH}_3\text{OH}$ or $[\text{Ni}(\text{C}_{10}\text{H}_{10}\text{N}_6)_2][\text{Ni}(\text{NCS})_3(\text{C}_{10}\text{H}_{10}\text{N}_6)_2] \cdot 2\text{CH}_3\text{OH}$. The asymmetric unit consists of half a centrosymmetric bis[tris(1*H*-pyrazol-1-yl)methane]-nickel(II) cation and an octahedral nickelate(II) anion bound to one tpm and three L ligands, and a methanol solvent molecule. One of the L ligands is disordered over two positions with occupancy factors of 0.650 (3) and 0.350 (3). There are $\text{O}-\text{H} \cdots \text{S}$ interactions between the methanol and the disordered thiocyanate anion, and a weak $\text{C}-\text{H} \cdots \text{O}$ hydrogen bond between the cation and the methanol O atom.

Related literature

For the ligand synthesis, see: Reger *et al.* (2000). For structural, spectroscopic and angular overlap studies of tris(1*H*-pyrazol-1-yl)methane complexes, see: Astley *et al.* (1993). For background information on the modelling of metallo-enzyme sites by small molecules, see: Kitajima *et al.* (1992); Trofimenko *et al.* (1992); Looney *et al.* (1992); Looney & Parkin (1994). A previous attempt to make similar building blocks with nickel(II) and a cyanide ligand is given in Lyubartseva & Parkin (2009).



Experimental

Crystal data

$[\text{Ni}(\text{C}_{10}\text{H}_{10}\text{N}_6)_2][\text{Ni}(\text{NCS})_3(\text{C}_{10}\text{H}_{10}\text{N}_6)_2] \cdot 2\text{CH}_3\text{O}$
 $M_r = 1445.65$
 Monoclinic, $C2/c$
 $a = 33.4463$ (8) Å
 $b = 7.3287$ (2) Å
 $c = 27.2689$ (7) Å

$\beta = 112.590$ (1)°
 $V = 6171.3$ (3) Å³
 $Z = 4$
 Cu $K\alpha$ radiation
 $\mu = 3.52$ mm⁻¹
 $T = 90$ K
 $0.20 \times 0.06 \times 0.02$ mm

Data collection

Bruker X8 Proteum diffractometer
 Absorption correction: multi-scan
 (SADABS in APEX2; Bruker, 2006)
 $T_{\text{min}} = 0.740$, $T_{\text{max}} = 0.933$

41685 measured reflections
 5605 independent reflections
 4932 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.061$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.128$
 $S = 1.11$
 5605 reflections
 418 parameters

6 restraints
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 1.34$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.44$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{O1S}-\text{H1S} \cdots \text{S3}^i$	0.84	2.43	3.267 (4)	175
$\text{O1S}-\text{H1S} \cdots \text{S3}^{ii}$	0.84	2.88	3.459 (6)	128
$\text{C23}-\text{H23} \cdots \text{O1S}$	1.00	2.15	3.118 (4)	162

Symmetry code: (i) $x - \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2006); cell refinement: SAINT (Bruker, 2006); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: XP in SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXL97 and local procedures.

GL gratefully acknowledges Southern Arkansas University for the financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG5255).

References

Astley, T., Gulbis, J. M., Hitchman, M. A. & Tiekink, E. R. T. (1993). *J. Chem. Soc. Dalton Trans.* pp. 509–515.

- Bruker (2006). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Kitajima, N., Fujisawa, K., Fujimoto, C., Morooka, Y., Hashimoto, S., Kitagawa, T., Toriumi, K., Tatsumi, K. & Nakamura, A. (1992). *J. Am. Chem. Soc.* **114**, 1277–1291.
- Looney, A. & Parkin, G. (1994). *Inorg. Chem.* **33**, 1234–1237.
- Looney, A., Parkin, G., Alsfasser, R., Ruf, M. & Vahrenkamp, H. (1992). *Angew. Chem. Int. Ed.* **31**, 92–93.
- Lyubartseva, G. & Parkin, S. (2009). *Acta Cryst.* **E65**, m1530.
- Reger, D. L., Grattan, T. C., Brown, K. J., Little, C. A., Lamba, J. J. S., Rheingold, A. L. & Sommer, R. D. (2000). *J. Organomet. Chem.* **607**, 120–128.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Trofimenko, S., Calabrese, J. C., Kochi, J. K., Wolowicz, S., Hulsbergen, F. B. & Reedijk, J. (1992). *Inorg. Chem.* **31**, 3943–3950.

supplementary materials

Acta Cryst. (2011). E67, m1656-m1657 [doi:10.1107/S1600536811045144]

Bis[tris(1*H*-pyrazol-1-yl- κ N²)methane]nickel(II) bis{[tris(1*H*-pyrazol-1-yl- κ N²)methane]tris(thiocyanato- κ N)nickelate(II)} methanol disolvate

G. Lyubartseva, S. Parkin and U. P. Mallik

Comment

Tripodal ligands with three pyrazolyl groups are increasingly being used in small-molecule modeling of the active sites of metallo-enzymes in which the metal is coordinated to two or three imidazole groups from histidine (Kitajima *et al.* 1992, Trofimenko *et al.* 1992, Looney *et al.* 1992, Looney & Parkin 1994). One of the goals of this research is to explore the chemistry of the neutral ligand tris(pyrazol-1-yl)methane compared to the more extensively studied isoelectronic but anionic ligand tris(pyrazol-1-yl)borate. In attempts to prepare mononuclear $[(\text{tpm})\text{Ni}^{\text{II}}L_3]^{-1}$, where tpm is tris(pyrazol-1-yl)methane, a symmetrical tripodal neutral nitrogen donor ligand, and L is NCS^- , a uninegative N-donor pseudohalide anion, we obtained $[(\text{tpm})_2\text{Ni}^{\text{II}}][(\text{tpm})\text{Ni}^{\text{II}}(\text{NCS})_3]_2 \cdot 2\text{CH}_3\text{OH}$ as blue monoclinic crystals in 57% isolated yield. The structure consists of centrosymmetric [bis[tris(1-pyrazolyl)methane- κ^3]-nickel(II) cations, with Ni^{II}—N distances ranging from 2.077 (2) to 2.082 (2) Å. The intraligand N—Ni—N angles in the cation range from 85.81 (10) to 95.27 (10)°, which introduces a slight trigonal distortion from perfect octahedral geometry. The anion consists of nickellate (II) atom surrounded octahedrally by one tripodal tris(pyrazol-1-yl)methane ligand and three isothiocyanate ligands. The tpm ligand N—Ni distances range from 2.080 (3) to 2.119 (3) Å, and the isothiocyanate N—Ni distances range from 2.046 (3) to 2.070 (3) Å.

Experimental

The tris(pyrazolyl)methane ligand was synthesized according to the previously published procedure by Reger *et al.* (2000). Tetrabutylammonium thiocyanate was purchased from Aldrich and used as received. $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ (475 mg, 2 mmol) was dissolved in 15 ml methanol. Tris(pyrazolyl) methane (428 mg, 2 mmol) was dissolved in 15 ml methanol. The ligand solution was added dropwise to metal solution and with moderate stirring. Once the addition was complete, tetrabutylammonium thiocyanate (1.81 g, 6 mmol) was added. The solution was filtered and methanol was evaporated slowly. Blue crystals were obtained after 3 days (549 mg, 57% yield). Elemental analysis, calculated for $\text{Ni}_3\text{C}_{48}\text{H}_{48}\text{N}_{30}\text{S}_6\text{O}_2$: C 39.88, H 3.35, N 29.07; found C 39.21, H 2.99, N 29.27%. IR (cm^{-1}): 3361, 3133, 2977, 2071, 1516, 1440, 1400, 1284, 1247, 1220, 1088, 1050, 980, 905, 858, 788, 766, 660, 608, 475.

Refinement

H atoms were found in difference Fourier maps and subsequently placed in idealized positions with constrained distances of 0.98 Å (RCH₃), 1.00 Å (R₃CH), 0.95 Å (C_{sp2}H), 0.84 Å (O—H), and with $U_{\text{iso}}(\text{H})$ values set to either $1.2U_{\text{eq}}$ or $1.5U_{\text{eq}}$ (RCH₃, OH) of the attached atom.

To ensure satisfactory refinement of disordered parts of the structure, a combination of constraints and restraints were used. The *SHELXL97* constraints EXYZ and EADP were used to make the geometry and displacement parameters of closely

supplementary materials

proximate disordered atoms equal. The *SHELXL97* restraint command DELU was also used to ensure similar displacement parameters for closely proximate, chemically similar groups.

The final weighting scheme (SHELXL-97 command "WGHT"), which is optimized to give a flat analysis of variance, had a somewhat larger than usual value for the second parameter. This is generally attributed to some form of bias, such as could be caused by unrecognized twinning or some other kind of incomplete model. We observed no obvious cause for the unusual weighting scheme, but the available sample was far from perfect. Indeed, the crystals were covered in a blue powder, which was likely caused by partial drying of the crystal. Some of this blue powder was easy to remove, but some was stuck to the crystal surface and could not be removed without damaging the crystals.

In the final difference map there are small residual peaks clustered around the disordered isothiocyanate sulphur. This could perhaps be due to partial occupancy/disordered solvent, but all attempts to model it as such did not improve the refinement enough to warrant retention of the extra details.

Figures




Fig. 1. View of the title compound with displacement ellipsoids drawn at the 50% probability level. The minor isothiocyanate disorder component is shown with open bonds. Unlabelled atoms are related to their labelled counterparts by inversion (0.5 - *x*, 1.5 - *y*, 1 - *z*).




Fig. 2. Packing diagram of the title compound as viewed down the *b* axis. The hydrogen atoms are omitted to enhance clarity.

Bis[tris(1*H*-pyrazol-1-yl- κ N²)methane]nickel(II) bis{[tris(1*H*-pyrazol-1-yl- κ N²)methane]tris(thiocyanato- κ N)nickelate(II)} methanol disolvate

Crystal data

$[\text{Ni}(\text{C}_{10}\text{H}_{10}\text{N}_6)_2][\text{Ni}(\text{NCS})_3(\text{C}_{10}\text{H}_{10}\text{N}_6)]_2 \cdot 2\text{CH}_4\text{O}$	$F(000) = 2968$
$M_r = 1445.65$	$D_x = 1.556 \text{ Mg m}^{-3}$
Monoclinic, <i>C2/c</i>	Cu <i>K</i> α radiation, $\lambda = 1.54178 \text{ \AA}$
Hall symbol: -C 2yc	Cell parameters from 9957 reflections
$a = 33.4463 (8) \text{ \AA}$	$\theta = 2.9\text{--}67.8^\circ$
$b = 7.3287 (2) \text{ \AA}$	$\mu = 3.52 \text{ mm}^{-1}$
$c = 27.2689 (7) \text{ \AA}$	$T = 90 \text{ K}$
$\beta = 112.590 (1)^\circ$	Rod, blue
$V = 6171.3 (3) \text{ \AA}^3$	$0.20 \times 0.06 \times 0.02 \text{ mm}$
$Z = 4$	

Data collection

Bruker X8 Proteum diffractometer	5605 independent reflections
Radiation source: fine-focus rotating anode graded multilayer optics	4932 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.061$
Detector resolution: 5.6 pixels mm^{-1}	$\theta_{\text{max}} = 68.0^\circ$, $\theta_{\text{min}} = 2.9^\circ$
φ and ω scans	$h = -39 \rightarrow 39$
Absorption correction: multi-scan (SADABS in APEX2; Bruker, 2006)	$k = -8 \rightarrow 8$
$T_{\text{min}} = 0.740$, $T_{\text{max}} = 0.933$	$l = -32 \rightarrow 32$
41685 measured reflections	

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.049$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.128$	H-atom parameters constrained
$S = 1.11$	$w = 1/[\sigma^2(F_o^2) + (0.0515P)^2 + 28.6001P]$
5605 reflections	where $P = (F_o^2 + 2F_c^2)/3$
418 parameters	$(\Delta/\sigma)_{\text{max}} = 0.001$
6 restraints	$\Delta\rho_{\text{max}} = 1.34 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.44 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. The crystals were covered in a blue powder, which was likely caused by partial drying of the crystal. Some of this blue stuff was easy to remove, but some was stuck to the crystal surface and could not be removed without damaging the crystal.

In the final difference map there are small residual peaks clustered around the disordered thiocyanate group. This could perhaps be due to partial occupancy/disordered solvent, but all attempts to model it as such did not improve the refinement enough to warrant retention of the extra details.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Ni1	0.388647 (18)	0.19922 (7)	0.82166 (2)	0.02134 (16)	
C1	0.38951 (11)	-0.2289 (4)	0.82452 (14)	0.0230 (7)	
H1	0.3895	-0.3653	0.8251	0.028*	
N1	0.33737 (9)	0.0180 (4)	0.80613 (11)	0.0212 (6)	

supplementary materials

N2	0.34542 (9)	-0.1642 (4)	0.80957 (11)	0.0196 (6)	
C2	0.29596 (11)	0.0339 (5)	0.79732 (13)	0.0221 (7)	
H2	0.2810	0.1467	0.7933	0.027*	
C3	0.27681 (11)	-0.1379 (5)	0.79473 (13)	0.0236 (7)	
H3	0.2474	-0.1630	0.7885	0.028*	
C4	0.30942 (11)	-0.2617 (5)	0.80312 (13)	0.0214 (7)	
H4	0.3072	-0.3908	0.8042	0.026*	
N3	0.41898 (9)	0.0209 (4)	0.88658 (12)	0.0234 (6)	
N4	0.41572 (9)	-0.1623 (4)	0.87707 (11)	0.0220 (6)	
C5	0.44066 (11)	0.0375 (5)	0.93835 (14)	0.0253 (7)	
H5	0.4477	0.1506	0.9567	0.030*	
C6	0.45189 (12)	-0.1344 (5)	0.96261 (15)	0.0322 (8)	
H6	0.4677	-0.1592	0.9992	0.039*	
C7	0.43519 (11)	-0.2583 (5)	0.92251 (15)	0.0290 (8)	
H7	0.4369	-0.3874	0.9258	0.035*	
N5	0.41142 (9)	0.0136 (4)	0.77985 (12)	0.0241 (6)	
N6	0.40755 (9)	-0.1675 (4)	0.78705 (12)	0.0226 (6)	
C8	0.43166 (12)	0.0246 (5)	0.74650 (15)	0.0286 (8)	
H8	0.4389	0.1359	0.7340	0.034*	
C9	0.44093 (12)	-0.1481 (5)	0.73221 (15)	0.0322 (8)	
H9	0.4551	-0.1762	0.7089	0.039*	
C10	0.42528 (11)	-0.2688 (5)	0.75883 (15)	0.0286 (8)	
H10	0.4266	-0.3982	0.7578	0.034*	
N7	0.36703 (10)	0.3584 (4)	0.86862 (12)	0.0280 (7)	
C11	0.36862 (11)	0.3894 (4)	0.91106 (15)	0.0253 (8)	
S1	0.37247 (3)	0.43823 (13)	0.97092 (4)	0.0346 (2)	
N8	0.35651 (10)	0.3522 (4)	0.75539 (12)	0.0276 (6)	
C12	0.33809 (11)	0.4581 (5)	0.72297 (13)	0.0221 (7)	
S2	0.31218 (3)	0.61217 (12)	0.67928 (3)	0.0267 (2)	
N9	0.44458 (11)	0.3525 (4)	0.84076 (15)	0.0371 (8)	0.650 (3)
C13	0.4780 (2)	0.3745 (8)	0.8323 (3)	0.0346 (11)	0.650 (3)
S3	0.52288 (6)	0.4092 (2)	0.82170 (10)	0.0554 (6)	0.650 (3)
N9'	0.44458 (11)	0.3525 (4)	0.84076 (15)	0.0371 (8)	0.350 (3)
C13'	0.4749 (4)	0.3881 (14)	0.8716 (5)	0.0346 (11)	0.350 (3)
S3'	0.52609 (11)	0.4382 (5)	0.91591 (17)	0.0639 (14)	0.350 (3)
Ni2	0.2500	0.7500	0.5000	0.01688 (18)	
N10	0.30477 (8)	0.5976 (4)	0.54331 (10)	0.0187 (6)	
N11	0.34180 (9)	0.6897 (4)	0.57058 (10)	0.0190 (6)	
C14	0.31541 (11)	0.4229 (4)	0.55193 (13)	0.0219 (7)	
H14	0.2959	0.3242	0.5379	0.026*	
C15	0.35916 (11)	0.4038 (5)	0.58445 (14)	0.0246 (7)	
H15	0.3746	0.2931	0.5964	0.030*	
C16	0.37506 (11)	0.5770 (4)	0.59556 (13)	0.0220 (7)	
H16	0.4040	0.6112	0.6168	0.026*	
N12	0.26851 (9)	0.9206 (4)	0.56604 (11)	0.0207 (6)	
N13	0.31151 (9)	0.9582 (3)	0.59102 (11)	0.0187 (6)	
C17	0.31975 (12)	1.0505 (5)	0.63671 (14)	0.0256 (7)	
H17	0.3473	1.0900	0.6611	0.031*	
C18	0.28065 (13)	1.0760 (5)	0.64097 (15)	0.0331 (9)	

H18	0.2755	1.1379	0.6686	0.040*
C19	0.24998 (12)	0.9929 (5)	0.59665 (14)	0.0262 (7)
H19	0.2198	0.9885	0.5893	0.031*
N14	0.20875 (8)	0.5902 (4)	0.52268 (11)	0.0195 (6)
N15	0.16941 (9)	0.5474 (4)	0.48424 (11)	0.0188 (6)
C20	0.20858 (11)	0.5150 (4)	0.56683 (13)	0.0220 (7)
H20	0.2319	0.5215	0.6004	0.026*
C21	0.16970 (12)	0.4255 (5)	0.55738 (14)	0.0262 (7)
H21	0.1618	0.3611	0.5826	0.031*
C22	0.14512 (11)	0.4486 (4)	0.50472 (13)	0.0217 (7)
H22	0.1166	0.4044	0.4860	0.026*
C23	0.15845 (11)	0.6126 (4)	0.43080 (13)	0.0188 (6)
H23	0.1287	0.5681	0.4084	0.023*
O1S	0.06346 (9)	0.4684 (4)	0.38493 (13)	0.0447 (7)
H1S	0.0533	0.3683	0.3705	0.067*
C1S	0.03005 (16)	0.5722 (8)	0.3923 (2)	0.0562 (13)
H1S1	0.0428	0.6501	0.4237	0.084*
H1S2	0.0090	0.4892	0.3974	0.084*
H1S3	0.0155	0.6484	0.3610	0.084*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ni1	0.0240 (3)	0.0130 (3)	0.0281 (3)	-0.0013 (2)	0.0112 (2)	0.0001 (2)
C1	0.0249 (17)	0.0149 (16)	0.0311 (18)	-0.0032 (13)	0.0129 (14)	-0.0018 (13)
N1	0.0254 (15)	0.0134 (13)	0.0269 (14)	0.0003 (11)	0.0123 (12)	0.0014 (11)
N2	0.0199 (14)	0.0145 (13)	0.0261 (14)	-0.0017 (10)	0.0107 (11)	-0.0015 (11)
C2	0.0239 (17)	0.0188 (16)	0.0255 (17)	0.0024 (13)	0.0114 (14)	0.0016 (13)
C3	0.0216 (17)	0.0243 (17)	0.0278 (17)	-0.0021 (14)	0.0126 (14)	0.0007 (14)
C4	0.0239 (17)	0.0174 (16)	0.0258 (16)	-0.0037 (13)	0.0128 (14)	-0.0021 (13)
N3	0.0241 (15)	0.0135 (13)	0.0336 (16)	-0.0005 (11)	0.0121 (12)	-0.0009 (11)
N4	0.0217 (14)	0.0152 (13)	0.0290 (15)	-0.0009 (11)	0.0093 (12)	-0.0010 (11)
C5	0.0193 (17)	0.0281 (18)	0.0273 (18)	-0.0020 (14)	0.0075 (14)	-0.0043 (14)
C6	0.0223 (18)	0.037 (2)	0.0326 (19)	-0.0025 (15)	0.0053 (15)	0.0040 (16)
C7	0.0219 (17)	0.0202 (17)	0.041 (2)	-0.0002 (14)	0.0079 (15)	0.0092 (15)
N5	0.0266 (15)	0.0175 (14)	0.0324 (16)	-0.0019 (11)	0.0158 (13)	0.0000 (12)
N6	0.0241 (15)	0.0153 (13)	0.0343 (16)	-0.0006 (11)	0.0176 (12)	-0.0031 (11)
C8	0.0275 (19)	0.0284 (19)	0.0330 (19)	-0.0053 (15)	0.0151 (15)	0.0003 (15)
C9	0.0298 (19)	0.036 (2)	0.037 (2)	-0.0051 (16)	0.0198 (16)	-0.0094 (17)
C10	0.0249 (18)	0.0234 (18)	0.040 (2)	-0.0006 (14)	0.0153 (16)	-0.0080 (15)
N7	0.0352 (17)	0.0172 (14)	0.0327 (17)	0.0006 (12)	0.0141 (13)	-0.0011 (12)
C11	0.0242 (18)	0.0132 (16)	0.041 (2)	-0.0018 (13)	0.0157 (15)	-0.0005 (14)
S1	0.0471 (6)	0.0277 (5)	0.0401 (5)	-0.0074 (4)	0.0290 (5)	-0.0065 (4)
N8	0.0303 (16)	0.0204 (15)	0.0350 (17)	-0.0018 (12)	0.0157 (13)	0.0014 (13)
C12	0.0238 (17)	0.0198 (17)	0.0264 (17)	-0.0048 (14)	0.0138 (14)	-0.0030 (14)
S2	0.0291 (5)	0.0257 (4)	0.0254 (4)	-0.0008 (3)	0.0104 (3)	0.0054 (3)
N9	0.0317 (18)	0.0188 (16)	0.057 (2)	-0.0024 (13)	0.0135 (16)	0.0059 (14)
C13	0.040 (3)	0.019 (2)	0.042 (3)	-0.003 (2)	0.013 (2)	0.002 (2)

supplementary materials

S3	0.0371 (10)	0.0350 (10)	0.0990 (16)	-0.0052 (7)	0.0315 (10)	0.0063 (9)
N9'	0.0317 (18)	0.0188 (16)	0.057 (2)	-0.0024 (13)	0.0135 (16)	0.0059 (14)
C13'	0.040 (3)	0.019 (2)	0.042 (3)	-0.003 (2)	0.013 (2)	0.002 (2)
S3'	0.0323 (18)	0.056 (2)	0.077 (3)	-0.0159 (15)	-0.0082 (16)	0.0269 (19)
Ni2	0.0191 (4)	0.0123 (4)	0.0213 (4)	0.0001 (3)	0.0100 (3)	0.0002 (3)
N10	0.0208 (14)	0.0133 (13)	0.0237 (14)	-0.0007 (10)	0.0105 (11)	-0.0002 (10)
N11	0.0226 (14)	0.0145 (13)	0.0213 (13)	-0.0015 (11)	0.0099 (11)	0.0003 (10)
C14	0.0272 (18)	0.0106 (15)	0.0311 (18)	0.0003 (13)	0.0147 (14)	0.0007 (13)
C15	0.0263 (18)	0.0159 (16)	0.0324 (18)	0.0057 (13)	0.0121 (15)	0.0019 (13)
C16	0.0200 (17)	0.0206 (17)	0.0256 (17)	0.0019 (13)	0.0089 (13)	0.0049 (13)
N12	0.0208 (14)	0.0164 (13)	0.0268 (15)	-0.0007 (11)	0.0112 (12)	-0.0019 (11)
N13	0.0212 (14)	0.0128 (13)	0.0237 (14)	0.0002 (10)	0.0105 (11)	-0.0009 (10)
C17	0.0332 (19)	0.0195 (17)	0.0246 (17)	-0.0043 (14)	0.0117 (15)	-0.0058 (13)
C18	0.044 (2)	0.0280 (19)	0.035 (2)	-0.0018 (16)	0.0233 (18)	-0.0105 (16)
C19	0.0296 (19)	0.0216 (17)	0.0336 (19)	0.0016 (14)	0.0190 (15)	-0.0036 (14)
N14	0.0184 (14)	0.0172 (13)	0.0223 (14)	0.0001 (11)	0.0071 (11)	0.0009 (11)
N15	0.0202 (14)	0.0151 (13)	0.0227 (14)	-0.0019 (10)	0.0101 (11)	0.0007 (10)
C20	0.0254 (18)	0.0208 (17)	0.0228 (16)	0.0017 (13)	0.0125 (14)	0.0017 (13)
C21	0.0312 (19)	0.0238 (18)	0.0287 (18)	-0.0015 (14)	0.0171 (15)	0.0044 (14)
C22	0.0242 (17)	0.0164 (16)	0.0278 (17)	-0.0034 (13)	0.0134 (14)	0.0024 (13)
C23	0.0230 (16)	0.0135 (15)	0.0219 (16)	0.0000 (12)	0.0109 (13)	0.0009 (12)
O1S	0.0299 (15)	0.0500 (19)	0.0543 (19)	-0.0059 (13)	0.0164 (13)	-0.0060 (15)
C1S	0.049 (3)	0.067 (3)	0.061 (3)	0.001 (2)	0.031 (2)	0.005 (3)

Geometric parameters (Å, °)

Ni1—N8	2.046 (3)	Ni2—N14 ⁱ	2.077 (3)
Ni1—N7	2.058 (3)	Ni2—N14	2.077 (3)
Ni1—N9	2.070 (3)	Ni2—N12 ⁱ	2.081 (3)
Ni1—N1	2.080 (3)	Ni2—N12	2.081 (3)
Ni1—N5	2.097 (3)	Ni2—N10 ⁱ	2.082 (3)
Ni1—N3	2.119 (3)	Ni2—N10	2.082 (3)
C1—N6	1.444 (4)	N10—C14	1.325 (4)
C1—N4	1.447 (4)	N10—N11	1.355 (4)
C1—N2	1.450 (4)	N11—C16	1.342 (4)
C1—H1	1.0000	N11—C23 ⁱ	1.449 (4)
N1—C2	1.317 (4)	C14—C15	1.396 (5)
N1—N2	1.358 (4)	C14—H14	0.9500
N2—C4	1.352 (4)	C15—C16	1.365 (5)
C2—C3	1.402 (5)	C15—H15	0.9500
C2—H2	0.9500	C16—H16	0.9500
C3—C4	1.369 (5)	N12—C19	1.327 (4)
C3—H3	0.9500	N12—N13	1.362 (4)
C4—H4	0.9500	N13—C17	1.349 (4)
N3—C5	1.322 (5)	N13—C23 ⁱ	1.447 (4)
N3—N4	1.364 (4)	C17—C18	1.369 (5)
N4—C7	1.355 (5)	C17—H17	0.9500
C5—C6	1.405 (5)	C18—C19	1.390 (5)

C5—H5	0.9500	C18—H18	0.9500
C6—C7	1.365 (6)	C19—H19	0.9500
C6—H6	0.9500	N14—C20	1.326 (4)
C7—H7	0.9500	N14—N15	1.367 (4)
N5—C8	1.328 (5)	N15—C22	1.358 (4)
N5—N6	1.355 (4)	N15—C23	1.441 (4)
N6—C10	1.359 (4)	C20—C21	1.389 (5)
C8—C9	1.394 (5)	C20—H20	0.9500
C8—H8	0.9500	C21—C22	1.364 (5)
C9—C10	1.369 (5)	C21—H21	0.9500
C9—H9	0.9500	C22—H22	0.9500
C10—H10	0.9500	C23—N13 ⁱ	1.447 (4)
N7—C11	1.160 (5)	C23—N11 ⁱ	1.449 (4)
C11—S1	1.627 (4)	C23—H23	1.0000
N8—C12	1.160 (5)	O1S—C1S	1.429 (6)
C12—S2	1.629 (4)	O1S—H1S	0.8400
N9—C13	1.237 (7)	C1S—H1S1	0.9800
C13—S3	1.652 (7)	C1S—H1S2	0.9800
C13'—S3'	1.713 (12)	C1S—H1S3	0.9800
N8—Ni1—N7	92.74 (12)	N14 ⁱ —Ni2—N12 ⁱ	95.27 (10)
N8—Ni1—N9	92.42 (13)	N14—Ni2—N12 ⁱ	84.73 (10)
N7—Ni1—N9	92.02 (14)	N14 ⁱ —Ni2—N12	84.73 (10)
N8—Ni1—N1	93.31 (12)	N14—Ni2—N12	95.27 (10)
N7—Ni1—N1	91.68 (12)	N12 ⁱ —Ni2—N12	179.996 (1)
N9—Ni1—N1	173.02 (12)	N14 ⁱ —Ni2—N10 ⁱ	93.96 (10)
N8—Ni1—N5	93.02 (12)	N14—Ni2—N10 ⁱ	86.04 (10)
N7—Ni1—N5	173.64 (12)	N12 ⁱ —Ni2—N10 ⁱ	85.81 (10)
N9—Ni1—N5	90.41 (13)	N12—Ni2—N10 ⁱ	94.19 (10)
N1—Ni1—N5	85.31 (11)	N14 ⁱ —Ni2—N10	86.04 (10)
N8—Ni1—N3	175.06 (12)	N14—Ni2—N10	93.96 (10)
N7—Ni1—N3	89.98 (12)	N12 ⁱ —Ni2—N10	94.19 (10)
N9—Ni1—N3	91.61 (12)	N12—Ni2—N10	85.81 (10)
N1—Ni1—N3	82.48 (11)	N10 ⁱ —Ni2—N10	180.00 (15)
N5—Ni1—N3	84.08 (11)	C14—N10—N11	104.9 (3)
N6—C1—N4	109.7 (3)	C14—N10—Ni2	137.4 (2)
N6—C1—N2	110.8 (3)	N11—N10—Ni2	117.63 (19)
N4—C1—N2	109.4 (3)	C16—N11—N10	112.2 (3)
N6—C1—H1	109.0	C16—N11—C23 ⁱ	128.6 (3)
N4—C1—H1	109.0	N10—N11—C23 ⁱ	119.3 (3)
N2—C1—H1	109.0	N10—C14—C15	110.7 (3)
C2—N1—N2	105.4 (3)	N10—C14—H14	124.6
C2—N1—Ni1	135.1 (2)	C15—C14—H14	124.6
N2—N1—Ni1	119.2 (2)	C16—C15—C14	105.8 (3)
C4—N2—N1	111.6 (3)	C16—C15—H15	127.1
C4—N2—C1	128.5 (3)	C14—C15—H15	127.1
N1—N2—C1	119.6 (3)	N11—C16—C15	106.4 (3)

supplementary materials

N1—C2—C3	111.0 (3)	N11—C16—H16	126.8
N1—C2—H2	124.5	C15—C16—H16	126.8
C3—C2—H2	124.5	C19—N12—N13	105.2 (3)
C4—C3—C2	105.5 (3)	C19—N12—Ni2	136.8 (2)
C4—C3—H3	127.2	N13—N12—Ni2	117.43 (19)
C2—C3—H3	127.2	C17—N13—N12	111.4 (3)
N2—C4—C3	106.5 (3)	C17—N13—C23 ⁱ	129.2 (3)
N2—C4—H4	126.8	N12—N13—C23 ⁱ	119.2 (3)
C3—C4—H4	126.8	N13—C17—C18	106.5 (3)
C5—N3—N4	105.3 (3)	N13—C17—H17	126.8
C5—N3—Ni1	136.5 (2)	C18—C17—H17	126.8
N4—N3—Ni1	118.1 (2)	C17—C18—C19	106.0 (3)
C7—N4—N3	111.3 (3)	C17—C18—H18	127.0
C7—N4—C1	128.7 (3)	C19—C18—H18	127.0
N3—N4—C1	119.6 (3)	N12—C19—C18	110.8 (3)
N3—C5—C6	111.0 (3)	N12—C19—H19	124.6
N3—C5—H5	124.5	C18—C19—H19	124.6
C6—C5—H5	124.5	C20—N14—N15	105.2 (3)
C7—C6—C5	105.4 (3)	C20—N14—Ni2	137.7 (2)
C7—C6—H6	127.3	N15—N14—Ni2	117.06 (19)
C5—C6—H6	127.3	C22—N15—N14	111.0 (3)
N4—C7—C6	107.0 (3)	C22—N15—C23	129.3 (3)
N4—C7—H7	126.5	N14—N15—C23	119.7 (3)
C6—C7—H7	126.5	N14—C20—C21	111.0 (3)
C8—N5—N6	105.1 (3)	N14—C20—H20	124.5
C8—N5—Ni1	136.0 (2)	C21—C20—H20	124.5
N6—N5—Ni1	118.8 (2)	C22—C21—C20	106.2 (3)
N5—N6—C10	111.5 (3)	C22—C21—H21	126.9
N5—N6—C1	119.8 (3)	C20—C21—H21	126.9
C10—N6—C1	128.5 (3)	N15—C22—C21	106.6 (3)
N5—C8—C9	111.2 (3)	N15—C22—H22	126.7
N5—C8—H8	124.4	C21—C22—H22	126.7
C9—C8—H8	124.4	N15—C23—N13 ⁱ	110.5 (3)
C10—C9—C8	105.5 (3)	N15—C23—N11 ⁱ	110.8 (3)
C10—C9—H9	127.2	N13 ⁱ —C23—N11 ⁱ	110.3 (3)
C8—C9—H9	127.2	N15—C23—H23	108.4
N6—C10—C9	106.6 (3)	N13 ⁱ —C23—H23	108.4
N6—C10—H10	126.7	N11 ⁱ —C23—H23	108.4
C9—C10—H10	126.7	C1S—O1S—H1S	109.5
C11—N7—Ni1	147.0 (3)	O1S—C1S—H1S1	109.5
N7—C11—S1	177.7 (3)	O1S—C1S—H1S2	109.5
C12—N8—Ni1	170.1 (3)	H1S1—C1S—H1S2	109.5
N8—C12—S2	177.7 (3)	O1S—C1S—H1S3	109.5
C13—N9—Ni1	144.9 (4)	H1S1—C1S—H1S3	109.5
N9—C13—S3	178.5 (5)	H1S2—C1S—H1S3	109.5
N14 ⁱ —Ni2—N14	180.00 (10)		
N8—Ni1—N1—C2	-54.5 (3)	N8—Ni1—N7—C11	-174.3 (5)

N7—Ni1—N1—C2	38.3 (3)	N9—Ni1—N7—C11	-81.8 (5)
N5—Ni1—N1—C2	-147.3 (3)	N1—Ni1—N7—C11	92.3 (5)
N3—Ni1—N1—C2	128.1 (3)	N3—Ni1—N7—C11	9.8 (5)
N8—Ni1—N1—N2	133.0 (2)	N8—Ni1—N9—C13	-83.5 (6)
N7—Ni1—N1—N2	-134.1 (2)	N7—Ni1—N9—C13	-176.4 (6)
N5—Ni1—N1—N2	40.2 (2)	N5—Ni1—N9—C13	9.5 (6)
N3—Ni1—N1—N2	-44.4 (2)	N3—Ni1—N9—C13	93.6 (6)
C2—N1—N2—C4	-0.2 (4)	N14 ⁱ —Ni2—N10—C14	-135.9 (3)
Ni1—N1—N2—C4	174.3 (2)	N14—Ni2—N10—C14	44.1 (3)
C2—N1—N2—C1	-173.8 (3)	N12 ⁱ —Ni2—N10—C14	-40.9 (3)
Ni1—N1—N2—C1	0.7 (4)	N12—Ni2—N10—C14	139.1 (3)
N6—C1—N2—C4	127.4 (3)	N14 ⁱ —Ni2—N10—N11	41.9 (2)
N4—C1—N2—C4	-111.5 (4)	N14—Ni2—N10—N11	-138.1 (2)
N6—C1—N2—N1	-60.1 (4)	N12 ⁱ —Ni2—N10—N11	136.9 (2)
N4—C1—N2—N1	61.0 (4)	N12—Ni2—N10—N11	-43.1 (2)
N2—N1—C2—C3	-0.3 (4)	C14—N10—N11—C16	0.3 (3)
Ni1—N1—C2—C3	-173.5 (2)	Ni2—N10—N11—C16	-178.1 (2)
N1—C2—C3—C4	0.6 (4)	C14—N10—N11—C23 ⁱ	179.5 (3)
N1—N2—C4—C3	0.5 (4)	Ni2—N10—N11—C23 ⁱ	1.0 (3)
C1—N2—C4—C3	173.5 (3)	N11—N10—C14—C15	-0.2 (4)
C2—C3—C4—N2	-0.7 (4)	Ni2—N10—C14—C15	177.8 (2)
N7—Ni1—N3—C5	-38.7 (3)	N10—C14—C15—C16	0.0 (4)
N9—Ni1—N3—C5	53.3 (4)	N10—N11—C16—C15	-0.3 (4)
N1—Ni1—N3—C5	-130.4 (3)	C23 ⁱ —N11—C16—C15	-179.4 (3)
N5—Ni1—N3—C5	143.5 (3)	C14—C15—C16—N11	0.1 (4)
N7—Ni1—N3—N4	137.9 (2)	N14 ⁱ —Ni2—N12—C19	144.3 (4)
N9—Ni1—N3—N4	-130.0 (2)	N14—Ni2—N12—C19	-35.7 (4)
N1—Ni1—N3—N4	46.2 (2)	N10 ⁱ —Ni2—N12—C19	50.7 (4)
N5—Ni1—N3—N4	-39.8 (2)	N10—Ni2—N12—C19	-129.3 (4)
C5—N3—N4—C7	0.0 (4)	N14 ⁱ —Ni2—N12—N13	-45.7 (2)
Ni1—N3—N4—C7	-177.6 (2)	N14—Ni2—N12—N13	134.3 (2)
C5—N3—N4—C1	173.4 (3)	N10 ⁱ —Ni2—N12—N13	-139.3 (2)
Ni1—N3—N4—C1	-4.3 (4)	N10—Ni2—N12—N13	40.7 (2)
N6—C1—N4—C7	-124.3 (4)	C19—N12—N13—C17	0.7 (4)
N2—C1—N4—C7	114.0 (4)	Ni2—N12—N13—C17	-172.2 (2)
N6—C1—N4—N3	63.7 (4)	C19—N12—N13—C23 ⁱ	176.2 (3)
N2—C1—N4—N3	-58.1 (4)	Ni2—N12—N13—C23 ⁱ	3.3 (3)
N4—N3—C5—C6	0.5 (4)	N12—N13—C17—C18	-1.0 (4)
Ni1—N3—C5—C6	177.4 (3)	C23 ⁱ —N13—C17—C18	-176.0 (3)
N3—C5—C6—C7	-0.8 (4)	N13—C17—C18—C19	0.9 (4)
N3—N4—C7—C6	-0.5 (4)	N13—N12—C19—C18	-0.1 (4)
C1—N4—C7—C6	-173.1 (3)	Ni2—N12—C19—C18	170.7 (3)
C5—C6—C7—N4	0.8 (4)	C17—C18—C19—N12	-0.5 (4)
N8—Ni1—N5—C8	50.3 (4)	N12 ⁱ —Ni2—N14—C20	138.3 (3)
N9—Ni1—N5—C8	-42.1 (4)	N12—Ni2—N14—C20	-41.7 (3)
N1—Ni1—N5—C8	143.4 (4)	N10 ⁱ —Ni2—N14—C20	-135.6 (3)

supplementary materials

N3—Ni1—N5—C8	-133.7 (4)	N10—Ni2—N14—C20	44.4 (3)
N8—Ni1—N5—N6	-132.6 (3)	N12 ⁱ —Ni2—N14—N15	-43.9 (2)
N9—Ni1—N5—N6	135.0 (3)	N12—Ni2—N14—N15	136.1 (2)
N1—Ni1—N5—N6	-39.5 (2)	N10 ⁱ —Ni2—N14—N15	42.2 (2)
N3—Ni1—N5—N6	43.4 (2)	N10—Ni2—N14—N15	-137.8 (2)
C8—N5—N6—C10	0.3 (4)	C20—N14—N15—C22	0.7 (3)
Ni1—N5—N6—C10	-177.6 (2)	Ni2—N14—N15—C22	-177.7 (2)
C8—N5—N6—C1	175.6 (3)	C20—N14—N15—C23	178.6 (3)
Ni1—N5—N6—C1	-2.3 (4)	Ni2—N14—N15—C23	0.2 (3)
N4—C1—N6—N5	-59.9 (4)	N15—N14—C20—C21	-0.3 (4)
N2—C1—N6—N5	60.9 (4)	Ni2—N14—C20—C21	177.6 (3)
N4—C1—N6—C10	114.5 (4)	N14—C20—C21—C22	-0.2 (4)
N2—C1—N6—C10	-124.6 (4)	N14—N15—C22—C21	-0.8 (4)
N6—N5—C8—C9	-0.1 (4)	C23—N15—C22—C21	-178.5 (3)
Ni1—N5—C8—C9	177.3 (3)	C20—C21—C22—N15	0.6 (4)
N5—C8—C9—C10	-0.1 (4)	C22—N15—C23—N13 ⁱ	-121.3 (3)
N5—N6—C10—C9	-0.4 (4)	N14—N15—C23—N13 ⁱ	61.3 (4)
C1—N6—C10—C9	-175.1 (3)	C22—N15—C23—N11 ⁱ	116.3 (4)
C8—C9—C10—N6	0.3 (4)	N14—N15—C23—N11 ⁱ	-61.2 (4)

Symmetry codes: (i) $-x+1/2, -y+3/2, -z+1$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1S—H1S \cdots S3 ⁱⁱ	0.84	2.43	3.267 (4)	175.
O1S—H1S \cdots S3 ⁱⁱⁱ	0.84	2.88	3.459 (6)	128.
C23—H23 \cdots O1S	1.00	2.15	3.118 (4)	162.

Symmetry codes: (ii) $x-1/2, -y+1/2, z-1/2$.

Fig. 1




Fig. 2

