

Acta Crystallographica Section D

**Biological
Crystallography**

ISSN 0907-4449

Crystallization and X-ray diffraction data for a new form of concanavalin A

Sean Parkin and George Craig

Copyright © International Union of Crystallography

Author(s) of this paper may load this reprint on their own web site provided that this cover page is retained. Republication of this article or its storage in electronic databases or the like is not permitted without prior permission in writing from the IUCr.

Crystallization and X-ray diffraction data for a new form of concanavalin A

Sean Parkin^{a*} and George Craig^b^aChemistry Department, University of Kentucky, Lexington, KY 40506, USA, and ^bLawrence Livermore National Laboratory, Livermore, CA 94550, USA

Correspondence e-mail: spark2@uky.edu

A new crystal form of concanavalin A, a well studied lectin from the jack bean (*Canavalia ensiformis*) is reported. These crystals, with the symmetry of orthorhombic space group $C222_1$, grow as large roughly equi-dimensional blocks. Unit-cell parameters at 120 K are $a = 118.67$ (12), $b = 101.36$ (13), $c = 111.94$ (9) Å. On density considerations for two molecules per asymmetric unit, the water content is ~60%. Data to a nominal resolution of 1.5 Å were collected.

Received 15 March 2002

Accepted 10 April 2002

1. Introduction

The structure of concanavalin A (conA) was first reported over 30 years ago (Reeke *et al.*, 1971; Hardman *et al.*, 1971; Edelman *et al.*, 1972; Hardman & Ainsworth, 1972) using orthorhombic ($I222$) crystals. The molecule consists of two large antiparallel β -sheets joined by numerous loops and turns. There are two native metal-binding sites, a Ca^{2+} site and a transition-metal site (predominantly Mn^{2+}), and there is evidence of a third (Naismith *et al.*, 1993). Selective repopulation of the transition-metal site markedly improves the diffraction limit of the $I222$ crystal form (Kalb *et al.*, 1988), allowing higher-resolution refinements to reveal ever greater detail (Emmerich *et al.*, 1994). True atomic resolution is attainable from native $I222$ conA crystals cooled to cryogenic temperatures, with concomitant increases in refinable detail (Parkin *et al.*, 1996; Deacon *et al.*, 1997). In spite of such high resolution, model accuracy and precision are difficult to assess in molecules as large and flexible as proteins (Kleywegt & Jones, 1997). Comparison of different crystal forms of the same species, however, can yield important information even at lower resolution. This provided the impetus to search for new crystal forms of this well studied protein.

2. Materials and methods

2.1. Crystallization

Crystals of conA (type 5V, C7275, Sigma-Aldrich, St Louis, MO, USA) grew in hanging drops containing 10 μ l of a 30 mg ml⁻¹ protein solution and 10 μ l precipitant [100 mM Tris-HCl pH 8.0 and 25%(v/v) ethanol] in water equilibrated against a 1 ml precipitant reservoir. Large block-shaped crystals formed over a few weeks.

2.2. Data collection

Data extending to 1.5 Å resolution were collected at 120 K on Xuong-Hamlin dual-multiwire area detectors attached to a Huber four-circle goniometer and Rigaku RU-200 Cu $K\alpha$ X-ray source. A crystal 0.4 mm on each side was washed briefly in antifreeze (~30% 2-methyl-2,4-pentanediol in the reservoir buffer), plunged into liquid nitrogen and mounted into the cold-stream path of a modified Siemens LT-2 low-temperature apparatus using specially designed tongs (Parkin & Hope, 1998).

3. Results

The crystals are orthorhombic, space group $C222_1$, with unit-cell parameters $a = 118.67$ (12), $b = 101.36$ (13), $c = 111.94$ (9) Å at 120 K. Given two molecules of conA per asymmetric unit, the solvent content is ~60%. Further details, including intensity statistics, appear in a separate paper (Kantardjieff *et al.*, 2002), where these crystals and data are used in a comparison with the two atomic resolution models from $I222$ form crystals (Parkin *et al.*, 1996; Deacon *et al.*, 1997).

We are grateful to the editor, to Peter Höchtel and to Rod Balhorn for valuable advice. This work was conducted at Lawrence Livermore National Laboratory under the auspices of the US Department of Energy, contract W-7405-ENG-48.

References

- Deacon, A., Gleichman, T., Kalb (Gilboa), A. J., Price, H., Raftery, J., Bradbrook, G., Yariv, J. & Helliwell, J. R. (1997). *J. Chem. Soc. Faraday Trans.* **93**, 4305–4312.
- Edelman, G. M., Cunningham, B. A., Reeke, G. N., Becker, J. W., Waxdal, M. J. & Wang, J. L. (1972). *Proc. Natl Acad. Sci. USA*, **69**, 2580–2584.
- Emmerich, C., Helliwell, J. R., Redshaw, M., Naismith, J. H., Harrop, S. J., Raftery, J., Kalb

- (Gilboa), A. J., Yariv, J., Dauter, Z. & Wilson, K. S. (1994). *Acta Cryst. D***50**, 749–756.
- Hardman, K. D. & Ainsworth, C. F. (1972). *Biochemistry*, **11**, 4910–4919.
- Hardman, K. D., Wood, M. K., Schiffer, M., Edmondson, A. B. & Ainsworth, C. F. (1971). *Cold Spring Harbor Symp. Quant. Biol.* **36**, 271–276.
- Kalb, A. J., Yariv, J., Helliwell, J. R. & Papiz, M. Z. (1988). *J. Cryst. Growth*, **88**, 537–540.
- Kantardjieff, K., Höchtel, P., Segelke, B., Tao, F.-M. & Rupp, B. (2002). *Acta Cryst. D***58**, 735–743.
- Kleywegt, G. J. & Jones, T. A. (1997). *Methods Enzymol.* **277**, 208–230.
- Naismith, J. H., Habash, J., Harrop, S., Helliwell, J. R., Hunter, W. N., Wan, T. C. M., Weisgerber, S., Kalb (Gilboa), A. J. & Yariv, J. (1993). *Acta Cryst. D***49**, 561–571.
- Parkin, S. & Hope, H. (1998). *J. Appl. Cryst.* **31**, 945–953.
- Parkin, S., Rupp, B. & Hope, H. (1996). *Acta Cryst. D***52**, 1161–1168.
- Reeke, G. N. Jr, Becker, J. W. & Quioco, F. A. (1971). *J. Biol. Chem.* **250**, 1525–1547.